


Review for Final

Shaoyun Yi

MATH 122

University of South Carolina

Spring 2021



- Average R.O.C. $\xrightarrow{\Delta t \rightarrow 0}$ R.O.C of $f(t)$ at $t = a$ (*Derivative* $f'(a)$).
- Accumulated Change $\xrightarrow{\Delta t \rightarrow 0}$ Definite Integral $\int_a^b f(t) dt$ (**F.T.C.**).
- https://people.math.sc.edu/shaoyun/Review_122F_SYi_Sp_21.pdf
- Composite Functions \leftrightarrow Chain Rule & Integration by Substitution

Functions

- Ask yourself:
- What is a function/domain (input)/range (output)?
 - Difference between **closed/open** intervals, i.e. **[,]** v.s. **(,)**
 - Interpret your numerical answers correctly
 - **Horizontal intercepts (Zeros)** & **Vertical intercept (0, f(0))**
 - Behavior: **increasing/decreasing/constant**
 - Concavity: Concave **up/down/neither (a line)**
- **Linear:** $y = mx + b$ & $y - y_0 = m(x - x_0)$.
 - **Exponential:** $P = P_0 \cdot a^t$ $\xleftrightarrow{\text{base change}}$ $P = P_0 \cdot e^{kt}$
 - P_0 : Initial quantity (Vertical intercept)
 - r : Decimal representation of percent rate of change ($r = a - 1$)
 - k : *Continuous* (growth/decay) rate of change
 - Base Change: $a = e^k \Leftrightarrow \ln a = k$
 - $a > 1$ (so $r > 0$ and $k > 0$): Exponential growth
 - $1 > a > 0$ (so $r < 0$ and $k < 0$): Exponential decay
 - **Natural Log.:** $y = \ln x \Leftrightarrow e^y = x$ ($x > 0$)
 - **Power (Poly.):** $Q(x) = k \cdot x^p$ ($P_n(x) = c_n x^n + \dots + c_1 x + c_0, c_n \neq 0$)
 - **Composite Functions:** $f(g(x))$. **Realize the inside function $u = g(x)$**

Changes

① Changes in y : $\Delta y = y_2 - y_1$. Unit: unit of y

② Average Rate of Change: Unit: unit of y per unit of t

$$\frac{\Delta y}{\Delta t} = \frac{f(b) - f(a)}{b - a} = \text{Slope of Secant line (between } a \text{ and } b)$$

- **Linear:** Its slope m is the constant Average Rate of Change.

- $\Delta t \rightarrow 0$: (Instantaneous) Rate of Change $f'(a)$ of f at a

$$f'(a) = \text{Slope of Tangent line at } A \text{ (or } x = a) \quad \dots (\star)$$

- Estimate $f'(a)$ by taking $\Delta t = 0.001$ in Average Rate of Change

>

- Determine $f'(a)$ is < 0 by reading the graph and using (\star)

=

- Estimate $f'(a)$ given numerically (Usually, Right-hand approximation)

- **First Derivative Test:** $f' > 0$ means $f \nearrow$ v.s. $f' < 0$ means $f \searrow$

③ Relative Change in P : $\frac{P_1 - P_0}{P_0}$ Unit: %

Applications

- **Average Rate of Change:** Average Velocity = $\frac{\text{change in distance}}{\text{change in time}}$
- **Linear:** Cost $C(q)$, Revenue $R(q)$, and Profit $\pi(q)$
 - $C(q) = C_{\text{fixed}} + C_{\text{variable}}$
 - $R(q) = p \cdot q$
 - $\pi(q) = R(q) - C(q)$
 - Break-even Point (Zero)
 - Marginal Cost/Revenue/Profit (Slopes \leftrightarrow Linear functions)
- **Exponential:** Know how to use “ $a = 1 + r$ ” and “ e^k ” properly
 - Exponential growth: has a constant Doubling Time
 - Exponential decay: has a constant Half-Life
 - Compound Interest:
$$\begin{cases} \text{annually} & P(t) = P_0 \cdot (1 + r)^t \\ \text{continuously} & P(t) = P_0 \cdot e^{rt} \end{cases}$$
- **Natural Log.:** Solve exponential equations using Natural Logarithm
- **Power:** Define a Polynomial (sum of power functions)
- **Composite Functions:** Produce New Functions from Old

How to Use Calculator

① Graph a Function:

(i) `y =`: (ii) `window`: X_{\min}/X_{\max} (iii) `zoom`: Choose “ZoomFit”

② Plot a Table of Data:

(i) `y =`: (ii) `stat`: **EDIT** & **1**: (iii) `zoom`: Choose “ZoomStat”

③ Find a Maximum:

(i) `y =`: (ii) `window`: X_{\min}/X_{\max} (iii) `zoom`: Choose “ZoomFit”
(iv) **2nd** + `trace`: Choose “maximum” (v) Left/RightBound?/Guess?

④ Evaluate a Value— $Y_1(X)$:

(i) `y =`: (ii) `vars`: **Y-VARS** **1**: **1**: Y_1 (iii) Main Screen: $Y_1(X)$

Formulas for Derivatives (*PDF is in Blackboard*)

- 1 **Constant Rule:** $(k)' = 0$
- 2 **Power Rule:** $(x^n)' = n x^{n-1}$
- 3 **Exponential Rule:** $(a^x)' = (\ln a) a^x$
- 4 **Exponential Rule:** $(e^x)' = e^x$
- 5 **Exponential Rule:** $(e^{kx})' = k e^{kx}$
- 6 **Natural Logarithmic Rule:** $(\ln x)' = \frac{1}{x}$
- 7 **Constant Multiple Rule:** $(c \cdot f)' = c \cdot f'$
- 8 **Sum Rule:** $(f + g)' = f' + g'$
- 9 **Difference Rule:** $(f - g)' = f' - g'$
- 10 **Product Rule:** $(f \cdot g)' = f' \cdot g + f \cdot g'$
- 11 **Quotient Rule:** $\left(\frac{f}{g}\right)' = \frac{f' \cdot g - f \cdot g'}{g^2}$
- 12 **Chain Rule:** $(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$

(Instantaneous) Rate of Change/Derivative

- 1 Leibniz's Notation: $f'(x) = \frac{dy}{dx}$ and $f'(a) = \frac{dy}{dx} \Big|_{x=a}$
- 2 Interpretations: **WWHHmU**
- 3 Tangent Line Approximation: $f(x) \approx f(a) + f'(a) \cdot (x - a)$
- 4 Relative rate of change of y at a is $\frac{f'(a)}{f(a)}$ and **Unit** is % per unit of t
- 5 **1st derivative test**: $f' > 0$ means $f \nearrow$ **v.s.** $f' < 0$ means $f \searrow$
- 6 **2nd derivative test**: $f'' > 0$ means $f \text{ 😊}$ **v.s.** $f'' < 0$ means $f \text{ 😞}$
- 7 **All derivative rules**: *C, Power/Exp/NLog, CM, S/D, Prod./Quot., Chain*
 - Find Equation of tangent line: $m = f'(a)$ & tangent point $(a, f(a))$
 - Find Second derivative: $f'' = (f')'$
 - Find Marginal Revenue/Cost/Profit: e.g. $MR = R'(q)$, MC , MP .
 - Find Specific values: Review *Examples in §3.3 and §3.4*
- 8 **Local Maxima/Minima**: Critical points ($f'(p) = 0$) & f' sign changes
Method: **2nd derivative test** ($f''(p) \neq 0$); **otherwise**, **1st derivative test**
- 9 **Global Maxima/Minima**: **Compare** critical values and endpoints values
Application in real life: Find Maximum/Minimum profit ($MP = \pi'(q) = 0$)
- 10 **Inflection points** (f changes concavity): $f''(p) = 0$ & f'' sign changes

Definite Integral

$$\begin{cases} \text{Left-hand sum} = f(t_0) \cdot \Delta t + f(t_1) \cdot \Delta t + \cdots + f(t_{n-1}) \cdot \Delta t \\ \text{Right-hand sum} = f(t_1) \cdot \Delta t + f(t_2) \cdot \Delta t + \cdots + f(t_n) \cdot \Delta t \end{cases}$$

$$\Delta t = \frac{b-a}{n}$$

Definite integral $\int_a^b f(t) dt := \lim_{n \rightarrow \infty} (\text{Left/Right-hand sum})$ (i.e., $\Delta t \rightarrow 0$)

• Use a calculator: **math** & choose "9." when you know f .

• **Geometric side:** • If $f(t) > 0$, then $\int_a^b f(t) dt =$ area under graph of f

• If $f(t) < 0$, then $\int_a^b f(t) dt = -$ area between a and b

• Area between two curves

• Estimate a definite integral *numerically/graphically*: **Left-/Right- sum**

• Interpretation: unit for $\int_a^b f(t) dt =$ **product** of unit of f and unit of t

• **The Fundamental Theorem of Calculus:** $\int_a^b F'(x) dx = F(b) - F(a)$

e.g. Total cost $C(b)$ of producing b units: $C(b) = C(0) + \int_0^b C'(q) dq$.

Indefinite Integral & Use FTC to evaluate definite integrals

- ① **Antiderivative:** If $F'(x) = f(x)$, then $F(x)$ is an *antiderivative* of $f(x)$.
Analyzing antiderivative graphically (FDT, FTC) & numerically (Calculator)
- ② **The Indefinite Integral of $f(x)$:** $\int f(x) dx = F(x) + C$
 - The *family* of antiderivatives of $f(x)$;
 - **Formulas for Antiderivatives:** $f(x) = k$; x^n ($n \neq -1$); $\frac{1}{x}$; e^x ; e^{kx} ;
 - Properties of Antiderivatives:** Sums & Constant Multiples
 - **Integration by substitution (u -substitution):** $u = u(x)$, $du = u'(x) dx$
- ③ Using **FTC** to evaluate Definite Integrals:
 - (i) Find an antiderivative $F(x)$, i.e., Calculate $\int f(x) dx = F(x) + C$
 - (ii) Evaluate the definite integral: $\int_a^b f(x) dx = F(x) \Big|_a^b = F(b) - F(a)$
 - (iii) A warm tip: *You can use a calculator to double-check your answers!*
 - (iv) Applications: • Find the area; • Evaluate definite integrals by **u -sub**.

i) **Constant Rule:** $(k)' = 0$

i) $\int k dx = kx + C$

ii) **Power Rule:** $(x^n)' = n x^{n-1}$

ii) $\int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1$

iii) **Exponential Rule:** $(a^x)' = (\ln a) a^x$

iii) $\int a^x dx = \frac{a^x}{\ln a} + C$

iv) **Exponential Rule:** $(e^x)' = e^x$

iv) $\int e^x dx = e^x + C$

v) **Exponential Rule:** $(e^{kx})' = k e^{kx}$

v) $\int e^{kx} dx = \frac{e^{kx}}{k} + C$

vi) **Natural Logarithmic Rule:** $(\ln x)' = \frac{1}{x}$

vi) $\int \frac{1}{x} dx = \ln|x| + C$

vii) **Constant Multiple Rule:** $(c \cdot f)' = c \cdot f'$

vii) $\int c \cdot f(x) dx = c \cdot \int f(x) dx$

viii) **Sum Rule:** $(f + g)' = f' + g'$

viii) $\int (f(x) + g(x)) dx = \int f(x) dx + \int g(x) dx$

ix) **Difference Rule:** $(f - g)' = f' - g'$

ix) $\int (f(x) - g(x)) dx = \int f(x) dx - \int g(x) dx$

x) **Product Rule:** $(f \cdot g)' = f' \cdot g + f \cdot g'$

x) NA

xi) **Quotient Rule:** $\left(\frac{f}{g}\right)' = \frac{f' \cdot g - f \cdot g'}{g^2}$

xi) NA

xii) **Chain Rule:** $(f \circ g)'(x) = f'(g(x)) \cdot g'(x)$

xii) **u-substitution:** $du = u'(x) dx$

Additional Suggestions

Review:

- 1 Your WileyPlus homework
- 2 Your class notes
- 3 Quizzes
- 4 Tests 1-3 Review Problems with Solutions in Blackboard
- 5 Lecture recordings in Blackboard

Contact:

- Me (virtual Office Hours or by e-mail)
- Your SI Leader

Good luck with all finals!