

# Final Exam

Exam Date: June 19th-20th (12:30 pm, Saturday–12:30 pm, Sunday)  
Exam Length: 150 minutes

---

- Please submit your work on Blackboard **before Sunday (6/20) 12:30 pm.**
  - You are required to submit your work as a single pdf.
  - Please make sure your handwriting is clear enough to read. Thanks.
  - **You will be allowed to use your notes and the book during the exam.**
  - **No collaborations are allowed.**
  - **No consulting any online sources is allowed.**
  - **No late work will be accepted.**
  - Total score: *150 points.*
- 

(0) [5 points] Write the following honors code with your full name at the end. I understand that it is the responsibility of every member of the Carolina community to uphold and maintain the University of South Carolina's Honor Code. As a Carolinian, I certify that I have neither given nor received unauthorized aid on this exam. **Full name**

(1) [30 points] True or False:

- $\mathbf{R}$  is a group under multiplication. **False: 0 has no inverse.**
- A cyclic group is always abelian. **True**
- $[9]_{35}$  is a unit in  $\mathbf{Z}_{35}$ . **True**
- If  $|G| = 31$ , then  $G$  must be isomorphic to  $\mathbf{Z}_{31}$ . **True**
- $A_n$  is a normal subgroup of  $S_n$ . **True**
- $D_4 \cong \mathbf{Z}_2 \times \mathbf{Z}_2 \times \mathbf{Z}_2$ . **False:  $D_4$  is nonabelian., while  $\mathbf{Z}_2 \times \mathbf{Z}_2 \times \mathbf{Z}_2$  is abelian.**
- The order of  $gH$  in  $G/H$  is the smallest positive integer  $n$  such that  $g^n = e$ . **False:  $\dots g^n \in H$ .**
- The product of an even number of disjoint cycles is an even permutation. **False:  $(12)(345)$  is odd.**
- $\mathbf{Z}_{10} \times \mathbf{Z}_{10} \cong \mathbf{Z}_5 \times \mathbf{Z}_{20}$ . **False: There is no element of order 20 in  $\mathbf{Z}_{10} \times \mathbf{Z}_{10}$ , while  $\mathbf{Z}_5 \times \mathbf{Z}_{20}$  does.**
- $\mathbf{Z}_{17}$  is a simple group. **True**

(2) [15 points] For the permutation  $\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 3 & 8 & 5 & 2 & 10 & 9 & 1 & 4 & 6 & 7 \end{pmatrix}$ :

(a) Write  $\sigma$  as a product of disjoint cycles.

$$\sigma = (135 \ 10 \ 7)(284)(69)$$

(b) Write  $\sigma$  as a product of transpositions.

$$\sigma = (13)(35)(5 \ 10)(10 \ 7)(28)(84)(69) = (10 \ 7)(57)(37)(17)(84)(24)(69)$$

(c) Write  $\sigma^{-1}$  as a product of disjoint cycles.

$$\sigma^{-1} = (17 \ 10 \ 53)(248)(69) \text{ since disjoint cycles commute.}$$

(d) Is  $\sigma$  even, odd, neither or both? Is  $\sigma^{-1}$  even, odd, neither or both?

odd

(e) What is the order of  $\sigma$ ?

$$\text{lcm}[5, 3, 2] = 30.$$

(3) [20 points] Let  $G$  be the set of nonzero rational numbers  $\mathbf{Q}^\times$ . Define a new multiplication by  $a * b = \frac{ab}{5}$ , for all  $a, b \in G$ . Show that  $(G, *)$  is an abelian group.

(i) Closure: Trivial since  $a, b \in \mathbf{Q}^\times$ .

(ii) Associative: For any  $a, b, c \in G$ , we have

$$(a * b) * c = \frac{ab}{5} * c = \frac{abc}{25} = a * \frac{bc}{5} = a * (b * c)$$

commutative:  $a * b = \frac{ab}{5} = \frac{ba}{5} = b * a$

(iii) Identity: The identity element is 5. For any  $a \in \mathbf{Q}^\times$ , we have  $5 * a = \frac{5a}{5} = a$ .

(iv) Inverses: For any  $a \in \mathbf{Q}^\times$ , its inverse is  $\frac{25}{a} \in \mathbf{Q}^\times$  since  $a \in \mathbf{Q}^\times$ . In fact,

$$a * \frac{25}{a} = \frac{a \cdot 25/a}{5} = 5.$$

For parts (iii)-(iv), we only check one equation because of the commutativity.

(4) (a) [5 points] Let  $G = \langle a \rangle$  be a group of order 50. What is the order of  $\langle a^{35} \rangle$ ?

The order of  $\langle a^{35} \rangle$  is  $\frac{50}{\gcd(35, 50)} = 10$ .

(b) [5 points] What is the order of  $([18]_{20}, [25]_{30})$  in  $\mathbf{Z}_{20} \times \mathbf{Z}_{30}$ ?

$o([18]_{20}) = \frac{20}{\gcd(18, 20)} = 10$  and  $o([25]_{30}) = \frac{30}{\gcd(25, 30)} = 6$ . Thus, we have

$$o(([18]_{20}, [25]_{30})) = \text{lcm}[10, 6] = 30.$$

(c) [5 points] Let  $G = \mathbf{Z}_{48}$ . List all possible choice of  $[k]_{48}$  such that  $\langle [k]_{48} \rangle = \langle [20]_{48} \rangle$ .

$\langle [k]_{48} \rangle = \langle [20]_{48} \rangle = \langle [4]_{48} \rangle \Rightarrow \gcd(k, 48) = 4 \Rightarrow \gcd(\frac{k}{4}, 12) = 1$ . And so all possible choice of  $[k]_{48}$  such that  $\langle [k]_{48} \rangle = \langle [20]_{48} \rangle$  are

$$[4]_{48}, [20]_{48}, [28]_{48}, [44]_{48}.$$

(5) [5 points] Let  $G$  be a non-cyclic group of order 27. Prove that  $a^9 = e$  for all  $a \in G$ .

*Proof.* Since  $G$  is not cyclic, it follows from Lagrange's theorem that an element  $a \in G$  can have order 1, 3 or 9. Hence proved.  $\square$

(6) Let  $H$  be a subgroup of  $G$ . Let  $N(H) = \{g \in G \mid gHg^{-1} = H\}$ . Prove

(a) [8 points]  $N(H)$  is a subgroup of  $G$ .

*Proof.*  $N(H)$  is nonempty since  $eHe^{-1} = H$ , i.e.,  $e \in N(H)$ .  
For any  $a, b \in N(H)$ , we have

$$abH(ab)^{-1} = a(bHb^{-1})a^{-1} = aHa^{-1} = H.$$

This implies that  $ab \in H$ . Finally, for any  $a \in N(H)$  we have

$$H = (a^{-1}a)H(a^{-1}a) = a^{-1}(aHa^{-1})a = a^{-1}H(a^{-1})^{-1} \text{ since } aHa^{-1} = H.$$

This implies that  $a^{-1} \in N(H)$ . □

(b) [6 points]  $H$  is a subgroup of  $N(H)$ .

*Proof.* It suffices to show that  $H \subseteq N(H)$  since both  $H$  and  $N(H)$  are subgroups of  $G$ . Thus, for any  $h \in H$ , we need to show  $hHh^{-1} = H$ .

$hHh^{-1} \subseteq H$  : For any  $h' \in H$ , we have  $hh'h^{-1} \in H$  since  $H$  is a subgroup.

$H \subseteq hHh^{-1}$  : For any  $h' \in H$ , we have  $h' = h(h^{-1}h'h)h^{-1} \in hHh^{-1}$  since  $h^{-1}h'h \in H$ . □

(c) [6 points]  $H$  is normal in  $N(H)$ .

*Proof.* For any  $h \in H$  and  $g \in N(H)$ , we have  $ghg^{-1} \in gHg^{-1} = H$ . □

(7) Let  $G$  and  $H$  be groups. Define the function  $\phi : G \times H \rightarrow G$  by

$$\phi((a, b)) = a, \text{ for all } (a, b) \in G \times H.$$

(a) [5 points] Prove that  $\phi$  is a group homomorphism and onto.

*Proof.* It is clear that  $\phi$  is well-defined and onto. For any  $(a_1, b_1), (a_2, b_2)$ , we have  $\phi((a_1, b_1), (a_2, b_2)) = \phi(a_1a_2, b_1b_2) = a_1a_2 = \phi((a_1, b_1))\phi((a_2, b_2))$ . □

(b) [5 points] Find  $\ker(\phi)$ .

$$\ker(\phi) = \{(a, b) \in G \times H \mid \phi((a, b)) = a = e_G\} = \{e_G\} \times H.$$

(8) (a) [4 points] Let  $G$  be an abelian group. Let  $H$  be a subgroup of  $G$ . Prove that  $aH = Ha$  for any  $a \in G$ .

*Proof.* Any subgroup of an abelian group is normal by commutativity. □

(b) [8 pts] List the cosets of  $\langle [11]_{24} \rangle$  in  $\mathbf{Z}_{24}^\times$ .

$$\mathbf{Z}_{24}^\times = \{[1]_{24}, [5]_{24}, [7]_{24}, [11]_{24}, [13]_{24}, [17]_{24}, [19]_{24}, [23]_{24}\}$$

$$\langle [11]_{24} \rangle = \{[1]_{24}, [11]_{24}\}.$$

$$[5]_{24}\langle [11]_{24} \rangle = \{[5]_{24}, [7]_{24}\}.$$

$$[13]_{24}\langle [11]_{24} \rangle = \{[13]_{24}, [23]_{24}\}.$$

$$[17]_{24}\langle [11]_{24} \rangle = \{[17]_{24}, [19]_{24}\}.$$

(c) [8 pts] Prove that the factor group  $\mathbf{Z}_{24}^\times / \langle [11]_{24} \rangle \cong \mathbf{Z}_2 \times \mathbf{Z}_2$ .

*Proof.* From part (b), we know that the factor group  $\mathbf{Z}_{24}^\times / \langle [11]_{24} \rangle$  has order 4. So it must be isomorphic to  $\mathbf{Z}_4$  or  $\mathbf{Z}_2 \times \mathbf{Z}_2$ . Moreover, every non-identity element in the factor group has order 2. In particular,  $[5]_{24}^2 = [1]_{24}$ ,  $[13]_{24}^2 = [1]_{24}$ , and  $[17]_{24}^2 = [1]_{24}$ . This implies that  $\mathbf{Z}_{24}^\times / \langle [11]_{24} \rangle \cong \mathbf{Z}_2 \times \mathbf{Z}_2$ . □

(9) [10 points] May you have a good summer! Stay safe!